

Human Health Hazard Assessment

Di-n-octyl Phthalate (DnOP)
(CAS No. 117-84-0)

TABLE OF CONTENTS

INTRODUCTION.....	3
1. IDENTITY	3
1.1 Identification of the Substance	3
1.2 Physico-Chemical Properties	3
2. USES	4
3. HUMAN HEALTH HAZARD	4
3.1 Toxicokinetics	4
3.2 Acute Toxicity	5
3.3 Irritation.....	5
3.4 Sensitisation	7
3.5 Repeated Dose Toxicity	7
3.6 Genetic Toxicity	10
3.7 Carcinogenicity	10
3.8 Reproductive Toxicity.....	11
3.8.1 Human Studies	11
3.8.2 Repeat Dose Toxicity Studies	12
3.8.3 Continuous Breeding Reproductive Toxicity Studies.....	12
3.8.4 Prenatal Developmental Toxicity Studies.....	13
3.8.5 Developmental/Postnatal Toxicity Studies	13
3.8.6 Mode of Action	13
4. HAZARD CHARACTERISATION	15
5. HUMAN HEALTH HAZARD SUMMARY TABLE.....	17
6. REFERENCES.....	18

INTRODUCTION

This review of Dioctyl phthalate (DnOP) is a health hazard assessment only. For this assessment, two key reviews on DnOP prepared by the Agency for Toxic Substances and Disease Registry (ATSDR, 1997) and the Centre for the Evaluation of Risks to Human Reproduction (CERHR, 2003) were consulted. These reviews were updated with relevant studies from more recent literature surveys conducted up to September 2006.


References not marked with an asterisk were examined for the purposes of this assessment. References not examined but quoted from these two reports as secondary citations are also noted in this assessment and marked with an asterisk.

Hazard information from this assessment is published also in the form of a phthalate hazard compendium providing a comparative analysis of key toxicity endpoints for 25 phthalates (NICNAS, 2007).

1. IDENTITY

1.1 Identification of the Substance

CAS Number: 117-84-0
 Chemical Name: 1,2-Benzenedicarboxylic acid, dioctyl ester
 Common Name: Dioctyl phthalate (DnOP)
 Molecular Formula: $C_{24}H_{38}O_4$
 Structural Formula:



Molecular Weight: 390.6
 Synonyms: Bis(n-octyl) phthalate; Di-n-octyl phthalate;; Dioctyl-o-benzenedicarboxylate; n-Dioctyl phthalate; n-Octyl phthalate; Octyl phthalate; Phthalic acid, dioctyl ester

Purity/Impurities/Additives None identified

1.2 Physico-Chemical Properties

Table 1: Summary of physico-chemical properties

<i>Property</i>	<i>Value</i>
Physical state	Colourless, odourless liquid
Melting point	-25°C
Boiling point	390°C (101.3 kPa)
Density	978 kg/m ³ (25°C)
Vapour pressure	1.91 x 10 ⁻⁵ kPa (25°C)
Water solubility	0.2, 3.0 x 10 ⁻³ g/L (25°C)
Partition coefficient n-octanol/water (log Kow)	5.22
Henry's law constant	0.55, 6.68 x 10 ⁻³ atm-m ³ /mole

Flash point 219°C
Source: ATSDR (1997)

2. USES

DnOP is manufactured in the United States predominantly as a minor component of C6-10 phthalate (approximately 20%). DnOP is used in this form in PVC for the manufacture of a variety of products including flooring and carpet tiles, canvas tarps, swimming pool liners, notebook covers, traffic cones, toys and dolls, vinyl furniture upholstery, shower curtains and gloves, garden hoses, weather stripping, flea collars and shoes. DnOP-containing PVC is also used in food applications such as seam cements, bottle cap liners and conveyor belts. Other non-PVC uses of DnOP include as a dye carrier in plastics production, for manufacture of adhesives, plastisols and nitrocellulose lacquer coatings, as an active pesticide ingredient and in cosmetics and colorants (ATSDR, 1997; CERHR, 2003).

In Australia, DnOP is imported for use as a plasticiser in automotive and industrial hose and for the manufacture of PVC conveyer belts, insulation materials, polyurethane surface coatings, floor finishes and adhesives. Imported toys, play and exercise balls contain DnOP. DnOP is also distributed to various institutions for research purposes.

3. HUMAN HEALTH HAZARD

3.1 Toxicokinetics

Previous Evaluations (ATSDR, 1997; CERHR, 2003)

Limited available data indicate that DnOP is readily absorbed following ingestion. Low levels of DnOP were reported in liver and adipose tissues of rats following administration in the diet over 13 weeks (Poon et al 1995*, 1997*). No data are available on absorption following inhalation or dermal administration.

Following oral administration of DnOP (2000 mg/kg) in rats, peak blood and testes levels of mono-*n*-octylphthalate (MnOP) were reached at 3 and 6 hours after dosing, respectively. Half-life of this metabolite in blood was 3.3 hours (Oishi 1990*). DnOP in addition to other dialkyl phthalates was reported to be metabolised to their monoesters and alcohol by enzymes present in gut tissues (Rowland et al., 1977*). Following administration of 0.2 mL of DnOP to rats by gavage, 31% of the administered dose was recovered in urine within 48 hours as both MnOP and free phthalic acid (Albro and Moore, 1974*).

Data not Reported in Previous Evaluations

In a more recent study of DnOP metabolism in rats, the monoester MnOP, mono-(3-caboxypropyl) phthalate (MCPP), phthalic acid and 5 additional oxidative products were detected in urine 24 hours after oral administration (300 mg/kg). MCPP was the major component with MnOP being the minor (Silva et al., 2005). In a similar study, MCPP levels were found to be 560-fold higher than MnOP levels in 24-hour urinary samples (Calafat et al., 2006). An *in vitro* study of the metabolism of DnOP by rat liver microsomes showed rapid production of MnOP with production also of mono-hydroxy-*n*-octyl phthalate and phthalic acid (Silva et al., 2005).

Conclusion

Following ingestion, DnOP is metabolised and rapidly absorbed from the gastrointestinal tract as the monoester mono-*n*-octylphthalate. Half-life of the monoester in the blood is approximately 3 hours. The liver is capable of metabolising DnOP. Elimination occurs via the urine with levels of MnOP exceeded after 24 hours by the other oxidative metabolite MCPP.

3.2 Acute Toxicity

Previous Evaluations (ATSDR, 1997; CERHR, 2003)

Results for acute toxicity are summarised in Table 2.

Table 2: Acute animal toxicity studies

<i>Study</i>	<i>Species</i>	<i>Result (LD₅₀)</i>	<i>Reference</i>
Oral	Rat	53700 mg/kg bw	Dogra et al (1987)*
Oral	Mouse	13000 mg/kg bw	Dogra et al (1989)*
Oral	Mouse	>12,800 mg/kg bw	Eastman Kodak Company (1978)*
Dermal	Guinea pig	75 ml/kg bw	Bisesi (1994); CMA (1999)*

Data not Reported in Previous Evaluations

No data.

Conclusion

DnOP has oral LD50 values of 53700 mg/kg bw in rats and >12,800 mg/kg bw in mice. A dermal LD50 value of 75 ml/kg bw in guinea pigs was reported, however, the concentration of the applied substances was not available. No data were available regarding inhalation toxicity.

3.3 Irritation

Skin Irritation

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997; CERHR, 2003)

DnOP was reported to be a slight skin irritant when applied to the depilated skin of guinea pigs (Eastman Kodak Company, 1978*). According to information from RTECS (2004), DnOP caused mild skin irritation when applied to rabbit skin (Marhold, 1986*).

Data on effects in humans are restricted to poorly documented clinical studies of skin irritation and sensitisation in volunteers upon dermal contact with dioctyl phthalate (isomers unspecified) (Government of Canada, 1993).

Data not Reported in Previous Evaluations

No data.

Conclusion

DnOP causes minimal skin irritation in guinea pigs. Limited data suggest similar effects in humans.

Eye Irritation

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997; RTECS 2004).

Ocular administration of DnOP to guinea pigs resulted in slight conjunctival irritation. No further details were provided (Eastman Kodak Company, 1978*).

According to information from RTECS (2004), application of 500 mg into the rabbit eye resulted in only mild eye irritation (Marhold, 1986*). Another poorly reported study in RTECS (authorship unknown) noted that application of DnOP (20 mg) into the rabbit eye caused severe eye irritation (Anon., 1946*).

Data on effects in humans are restricted to a case report of irritation of the eye of workers exposed to phthalates including dioctyl phthalate (isomer unspecified) (Government of Canada, 1993).

Data not Reported in Previous Evaluations

No data.

Conclusion

DnOP causes minimal eye irritation in guinea pigs. Limited data suggest similar effects in humans.

Respiratory Irritation

Previous Evaluations (Government of Canada, 1993)

Data are restricted to a case report of irritation of the upper respiratory tract of workers exposed to phthalates including dioctyl phthalate (isomer unspecified) (Government of Canada, 1993).

Data not Reported in Previous Evaluations

No data.

Conclusion

Data are insufficient to determine the respiratory irritation potential of DnOP.

3.4 Sensitisation

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997)

DnOP was not a skin sensitiser in guinea pigs (Eastman Kodak Company, 1978*). Data on effects in humans are restricted to poorly documented clinical studies of skin irritation and sensitisation in volunteers upon dermal contact with dioctyl phthalate (isomers unspecified).

Data not Reported in Previous Evaluations

A European Chemicals Bureau (ECB) IUCLID dataset for a related phthalate mixture likely containing DnOP (1,2-benzenedicarboxylic acid, di-C8-10-alkyl esters, CAS no. 71662-46-9) noted negative results from a guinea pig maximisation test conducted to OECD Test Guidelines and Good Laboratory Practice guidelines.

The ECB IUCLID dataset above also contains short summaries of human sensitisation data for DnOP. Sixty workers chosen from 400 workers in a plastic shoe factor were patch tested with a standard battery of substances including 4 phthalates. Of 11 workers with contact dermatitis, 6 workers showed sensitisation reactions. Reactions to DnOP were observed. In another summary of a case report of occupational asthma, DnOP was reported to evoke an asthmatic reaction in a single patient following exposure to DnOP vapours. The reaction was inhibited by prior administration of sodium cromoglycate.

No other details were provided for either of these summaries and the data have not undergone any evaluation by the European Commission (ECB, 2000).

Conclusion

A single animal study indicated that DnOP is not a skin sensitiser. Negative animal results were also noted above for a di-C8-10-alkyl ester mixture likely containing DnOP. In contrast, poorly documented human patch test studies and a case report from occupational exposure suggest the potential for skin and respiratory sensitisation in humans.

Overall, data are insufficient to determine the sensitisation potential of DnOP.

3.5 Repeated Dose Toxicity

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997; CERHR, 2003 and Ontario Ministry of the Environment, 2005).

Administration of DnOP (2% in diet) to male JCL:Wistar rats for 1 week produced statistically significant increases in absolute and relative liver weights but not kidney weights (Oishi and Hiraga, 1980). Administration of dietary mono-octyl phthalate also induced liver enlargement but also altered serum lipid composition (Oishi and Hiraga 1982*).

Wistar rats received DnOP orally at doses of 2266 mg/kg bw/d for 3 days, 2078 mg/kg bw/d for 10 days, and 1906 mg/kg bw/d for 21 days. The livers of treated rats had a pale greasy appearance. Relative liver weights were statistically significantly increased after 10 and 21 days of treatment, but not after 3 days of treatment. After 3 days, loss of centrilobular glycogen, proliferation and dilation of smooth endoplasmic reticulum, loss of rough endoplasmic reticulum, and shortened microvilli in some bile canaliculi were observed. No

significant changes were noted in enzyme parameters associated with peroxisome proliferation. After 10- or 21-day treatment, centrilobular glycogen reduction became more severe and was associated with fat accumulation and some necrosis. Small but statistically significant increases were also noted for hepatic cyanide-insensitive palmitoyl CoA oxidase and peroxisomal catalase activities for these prolonged treatments (Mann et al., 1985*).

Thyroid effects were re-examined in the above study. Serum from 4 Wistar rats administered DnOP for 21 days in the above study were re-assayed (Hinton et al., 1986*). Statistically significant decreases in serum thyroxine (T4) levels, but not triiodothyronine (T3) levels, were observed. In addition, in the thyroid, ultrastructural changes consisting of increases in the numbers and sizes of lysosomes, enlargement of the Golgi apparatus and damage to mitochondria were also detected.

To examine whether liver metabolism and biochemical changes are associated with peroxisome proliferation, 6 Sprague-Dawley rats/dose were treated by gavage with 1000 or 2000 mg/kg bw/d DnOP for 14 days. Relative liver weights were increased in both groups. There were no increases in peroxisomal enzyme activities (Lake et al., 1984*; 1986*).

In an incomplete report of a study in which male rats were exposed to 0, 300 or 600 mg/kg bw/d of DnOP in the diet for 11 weeks, marked liver damage (characterized by cellular enlargement and proliferation, vacuolization, chronic inflammation, and necrosis) was observed at 300 mg/kg bw/d and above. Alterations in serum enzyme levels were consistent with the observed liver damage (DeAngelo et al., 1988*).

Rats received DnOP intraperitoneally at doses of 100, 300, or 600 mg/kg bw/d for 90 days. Half of the animals from each dose group were killed at the end of the study. Microscopic examination of the kidneys revealed dose related changes mainly to glomeruli, proximal and distal tubules, the loop of Henle and collecting tubules in the medulla. Effects persisted 45 days after cessation of treatment (Khanna et al., 1990*).

Systemic effects of DnOP were studied in Sprague-Dawley rats (10/sex) fed DnOP at dietary concentrations of 0, 5, 50, 500, or 5,000 ppm (m/f: 0/0, 0.4/0.4, 3.5/4.1, 36.8/40.8, or 350/403 mg/kg bw/d) for 13 weeks. At 5000 ppm, liver and thyroid effects were observed. Hepatic effects including anisokaryosis, nuclear hyperchromicity, vesiculation, cytoplasmic vacuolation, nuclear endothelial prominence, and accentuation of zonation were observed. Statistically significant increases in hepatic ethoxyresorufin-o-demethylase activity were also seen at this dose level. Thyroid effects included decreases in follicle size and colloid density. There were no morphological effects on reproductive organs at the highest dose level. The NOAEL was 500 ppm (37 mg/kg bw/d), based on liver and thyroid effects at 5000 ppm (Poon et al., 1995*; 1997*).

DnOP was also studied in mice. Following a continuous breeding protocol, offspring CD-1 mice taken from the last litter of treated parental mice exposed to approximately 7500 mg/kg bw/d DnOP for 105 days, including mating, were exposed to DnOP for 85 to 105 days, first via lactation, then through diet. The average DnOP dose in these offspring over this period was calculated by the study authors as 8640 mg/kg bw/d. Statistically significant increases in absolute liver weights were observed in both sexes. Absolute kidney weights were also statistically significantly increased in females but not males. However, no gross morphological or histopathological lesions were noted in these two organs (Heindel et al, 1989, Morrissey et al., 1989*, NTP 1985*).

CD-1 mice receiving DnOP in the diet at concentrations of 1800, 3600, 7500, and 15000 mg/kg bw/d for 14 days showed no clinical symptoms of toxicity except for rough hair coats at the highest dose (Heindel et al, 1989*).

Information is limited on chronic toxicity of DnOP in experimental animals. In an incomplete account, a single dose (175 mg/kg bw/d) of an unspecified isomer of dioctyl phthalate was administered to an unspecified species and strain of animal for 12 months. The reported effects were restricted to a decrease in body weights and increases in liver and kidney weights and activities of serum enzymes (Pieckacz, 1971*). In another chronic toxicity study, "numerous" liver nodules were reported in male rats given 1% DnOP (600 mg/kg bw/d) in the diet for 15 months. The activity of a number of lysosomal enzymes was also increased in these rats (Carter et al., 1989*).

There are few human data for repeat dose exposure to DnOP. Government of Canada (1993) and Ontario Ministry of the Environment (2005) contain the following references to poorly documented studies of occupational exposure to phthalates including DnOP.

Gilioli et al. (1978*) reported that 12 of 23 workers at a plasticizer manufacturing facility who had been exposed to phthalate esters (< 1 to 60 mg/m³, specific isomers not identified) for an average of 4.5 years had a mild to moderate sensory-motor and motor polyneuropathy. Another study investigated Russian workers (87 females and 60 males) in the artificial leather industry in which several phthalate plasticisers including DnOP were used (Milkov et al., 1973*). Duration of employment ranged from 0.5 to 19 years. Phthalates were found to be the principal air contaminants in the work areas with ambient air concentrations of the phthalate plasticisers ranging from 1.7 to 66 mg/m³. Neurological testing revealed that 32% of the employees had polyneuritis, the incidence of which correlated positively with length of service. A subset (81 workers) was examined for vestibular function effects, with 78% of subjects showing a depression of vestibular receptors. Some workers also displayed a lowering of the excitability threshold level for olfactory receptors.

Data not Reported in Previous Evaluations

In a US New Jersey Department of Health and Senior Services Hazardous Substances Fact Sheet, repeated exposure of humans to DnOP is claimed to cause liver damage, while repeated skin contact can cause dryness, cracking and rashes (NJDHSS, 2002).

Conclusion

The liver appears to be the primary target organ of DnOP. Liver toxicity (weight, histological or clinical chemistry changes) caused by DnOP was observed in several repeat dose studies. In a 21-day oral study, DnOP caused dose-related statistically significant increases in relative liver weights, loss of centrilobular glycogen and changes in liver microarchitecture including fat accumulation and necrosis. In other studies, DnOP also induced ultrastructural changes in the thyroid and kidney.

Overall, liver effects from DnOP did not appear to be associated with peroxisome proliferation. Increases in peroxisomal enzyme activities generally were not observed.

From a 13-week rat dietary study of 4 dose levels, a LOAEL of 350 mg/kg bw/d was established, based on histological changes in the liver (anisokaryosis, nuclear hyperchromicity, vesiculation, cytoplasmic vacuolation, nuclear endothelial prominence, accentuation of zonation) and thyroid (decreases in follicle size and colloid density). The NOAEL was 37 mg/kg bw/d.

3.6 Genetic Toxicity

Previous Evaluations (ATSDR, 1997; CERHR, 2003).

ATSDR (1997) notes a sizable database of *in vitro* microbial assays for DnOP. DnOP ranging from 100 to 10000 µg/plate in either the presence or absence of exogenous metabolic activation was not mutagenic in Ames tests using *S. typhimurium* TA 1535, TA 1537, TA 98, TA 100 (Zeiger et al., 1982*; 1985*).

DnOP at 2000 µg/mL with metabolic activation and ranging from 100 to 2000 µg/mL without metabolic activation did not induce DNA damage in *E. coli* (Goodyear Tire & Rubber Company, 1981*). Similar results are reported by Seed (1982) and Shibamoto and Wei (1986*). DnOP also showed a negative response in a prokaryotic SOS chromotest assay to detect DNA damage in *E. coli* (Sato et al., 1994*).

Data on mixtures containing DnOP were noted in CERHR (2003). Results from testing a C6-10 phthalate mixture (containing 20% DnOP) in a mouse lymphoma mutation assay were considered equivocal due to a non-dose related increase in mutations in the presence and absence of metabolic activation (Barber et al., 2000). According to CMA (1999*), di(n-octyl, n-decyl) phthalate containing DnOP as a component was reported to be negative in an Ames test and the Chinese hamster ovary/HPRT locus assay.

Data not Reported in Previous Evaluations

No data.

Conclusion

DnOP is negative in bacterial mutation and DNA damage assays. No information is available regarding genotoxic potential *in vivo*.

Based on negative results from *in vitro* bacterial mutation and DNA damage assays, DnOP is considered non-genotoxic.

3.7 Carcinogenicity

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997; Ontario Ministry of Environment, 2005)

In two studies, male Sprague-Dawley rats were given a partial hepatectomy and then injected intraperitoneally with diethylnitrosamine, then exposed to 0.5 or 1% DnOP via the diet (approximately 250 mg/kg bw/d or 500 mg/kg bw/d) for up to 26 weeks. Increases in numbers of gamma-glutamyl transpeptidase-positive foci and levels of glutathione-S-transferase staining in the livers compared to controls was observed, suggesting pre-tumour activity. Only a 3-fold increase was observed for carnitine acetyltransferase activity, a marker for

peroxisome proliferation (DeAngelo et al., 1986*; 1989*; Carter et al., 1992). These results suggest that DnOP may promote preneoplastic lesions in rat livers via a mechanism that does not rely solely on peroxisome proliferation.

In a poorly documented chronic toxicity study, "numerous" liver nodules were reported in male rats given 1% DnOP (600 mg/kg bw/d) in the diet for 15 months. The activity of a number of lysosomal enzymes was also increased in these rats (Carter et al., 1989*).

Data from an *in vitro* test where a C6-10 phthalate mixture (containing 20% DnOP) was examined in the Balb/3T3 mammalian cell transformation assay showed a negative result (Barber et al., 2000).

Data not Reported in Previous Evaluations

No data.

Conclusion

Limited data suggest that dietary exposure to DnOP might act as a promoter of pre-neoplastic lesions in the rat liver. Overall, data are insufficient to determine the carcinogenic potential of DnOP.

3.8 Reproductive Toxicity

Traditional hazard assessments consider reproductive toxicity separate from developmental toxicity. Reproductive toxicity is tested by exposing sexually mature adults to a chemical and examining the effects on the animal's capacity to reproduce. Developmental toxicity is studied by exposing pregnant dams and looking for effects in the fetuses. However, these tests generally do not detect chemicals that induce effects that only manifest postnatally. Thus, chemicals that affect the developing reproductive system following prenatal exposure may also affect sexual maturation or functional reproductive disorders that are only apparent at maturity. Developmental toxicity can therefore lead to reproductive toxicity and the two endpoints cannot be clearly distinguished.

In this hazard assessment, data will be presented on the basis of test procedure. The effects on fertility and development will then be discussed separately in the conclusion.

3.8.1 Human Studies

Previous Evaluations (ATSDR, 1997; CERHR, 2003)

Human sperm suspensions incubated with DnOP (0, 64, 640 μ M) for up to 18 hours showed dose-dependent decreases in motility (80% motility at 64 μ M) (Fredricsson et al., 1993).

Data not Reported in Previous Evaluations

A group of 45 women with endometriosis showed significantly higher plasma concentrations of DnOP as well as DBP, BBP, DEHP and polychlorinated biphenyls compared to a group of 135 fertile women (Reddy et al., 2006a; b).

Associations between 11 maternal urinary phthalate monoester concentrations and genital parameters such as anogenital distance and testicular descent in children were studied in 85 mother-son pairs (Swan et al., 2005). No significant association was found between maternal urinary MCPP (a DnOP metabolite) concentration and infant anogenital index (AGI).

3.8.2 Repeat Dose Toxicity Studies

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997; CERHR, 2003 and Ontario Ministry of the Environment, 2005)

Male Sprague-Dawley rats receiving DnOP by gavage at 2800 mg/kg bw/d for 4 or 10 days showed no testicular atrophy or histological lesions, testicular zinc loss or prostate or seminal vesicle weight loss (Foster et al., 1980; Gray and Butterworth 1980*).

Similarly, no effects on testes weights or testicular concentration of testosterone or dihydrotestosterone were observed in male rats fed a diet containing 2% DnOP for one week. However, a statistically significant 15% reduction in testicular zinc was observed (Oishi & Hiraga, 1980). No effect on testis weight, gross morphology, or histopathology was found in male rats receiving dietary exposure to approximately 2,000 mg/kg/d for 10 or 21 days (Mann et al., 1985).

In a subchronic study, Sprague-Dawley rats received DnOP in diet at doses as high as 5000 ppm (m/f: 350/403 mg/kg bw/d) for 13 weeks. Testes weights were unaffected and testes histology was normal. No reproductive effects were seen (Poon et al., 1995*; 1997*).

Data not Reported in Previous Evaluations

No data.

3.8.3 Continuous Breeding Reproductive Toxicity Studies

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997; CERHR, 2003 and Ontario Ministry of the Environment, 2005)

In a continuous breeding protocol oral reproductive/developmental study, CD-1 Swiss mice (20 pairs/dose) were fed DnOP in the diet at 0, 1.25, 2.5, or 5% (0, 1800, 3600, and 7500 mg/kg bw/d) for 14 weeks (Gulati et al., 1985*; Heindel et al., 1989). Litters born during the 14 weeks period were evaluated and removed so that the adults could continue breeding. Over 5 successive litters, there was no effect on the number of litters, litter size, sex ratios, pup weight or viability.

In sexually mature F1 adults mated within dose groups, DnOP similarly had no effects on fertility indices, litter size, pup weight or viability. In control and high-dose (7500 mg/kg bw/d) F1 adults, increased absolute and relative liver weights and decreased relative seminal vesicles weights were observed in males. However, there was no evidence of morphological or histopathological changes in these organs. No changes were observed in testis, cauda epididymis or prostate weights, sperm concentrations, percent mobile sperm or percent abnormal sperm. Females at this high dose showed increased relative liver and kidney weights. However, no changes were observed in oestrous cycle length or in reproductive organ weights or histopathology. Over 5 litters, there was no effect on the number of litters, litter size, sex ratios, pup weight or viability. Overall, DnOP did not show significant

reproductive effects in this study even at hepatotoxic doses. The NOAEL for reproductive and developmental effects was 7500 mg/kg bw/d, the highest dose tested.

Data not Reported in Previous Evaluations

No data.

3.8.4 Prenatal Developmental Toxicity Studies

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997; CERHR, 2003 and Ontario Ministry of the Environment, 2005)

Singh et al. (1972) administered DnOP at 0, 4890, 9780 mg/kg bw/d to Sprague-Dawley rats (15/group) prenatally on gestation days 5, 10 and 15 via intraperitoneal injection. A small but significant decrease in average foetal weight and a significant increase in the incidence in gross foetal malformations were observed in offspring. Possible confounding maternal effects were not reported.

Administration of n-octanol, a primary metabolite of DnOP (~130, 650, 945, and 1,300 mg/kg bw/d) by gavage to pregnant Wistar rats on days 6–15 of gestation induced dose-related symptoms of clinical intoxication of the nervous system with maternal death seen in the three highest dose levels. However, no effects on fetal weight, viability, or incidence of malformations were seen (Hellwig and Jack, 1997*)

Data not Reported in Previous Evaluations

No data.

3.8.5 Developmental/Postnatal Toxicity Studies

Previous Evaluations (Government of Canada, 1993; ATSDR, 1997; CERHR, 2003 and Ontario Ministry of the Environment, 2005)

Hardin et al. (1987) administered DnOP via oral gavage to female CD-1 mice at 9780 mg/kg bw/d during GD 6-13. There was no effect on maternal weight. The test group varied from the concurrent control (vehicle treated) group only in an 11% reduction in numbers of liveborn pups per litter and a 14% reduction in average pup weight gains on postnatal days 1-3 (both statistically significant). The authors reported that concurrent control values for these parameters were higher than other control groups from the same study, thus casting uncertainty on the biological significance of these changes.

Data not Reported in Previous Evaluations

No data.

3.8.6 Mode of Action

Previous Evaluations (CERHR, 2003)

DnOP did not induce estrogenic responses in vivo in a uterotrophic and vaginal cornification assay using immature and mature ovariectomised rats (Zacharewski et al., 1998). DnOP was

negative for estrogenic activity in recombinant yeast assay (Harris et al., 1997) and was not a competitive agonist at the oestrogen receptor in an *in vitro* competitive ligand-binding assay and did not induce oestrogen receptor-mediated gene expression in MCF-7 cells (Zacharewski et al., 1998).

Phthalate-induced germ cell detachment was examined *in vitro* in co-cultures of Sertoli and germ cells isolated from pubertal rats. (Gray and Beamand, 1984). Mono-n-octyl phthalate (MnOP) induced marked germ-cell detachment and disruption of the Sertoli-cell monolayer at 10^{-4} M, the magnitude of which exceeded that of the other 8 monoesters tested.

Data not Reported in Previous Evaluations

DnOP (up to 10^{-3} M) showed no detectable binding to estrogen receptor α or β *in vitro* (Toda et al., 2004) and did not demonstrate estrogenic activities in *in vitro* reporter gene assays with CHO-K1 cells transfected with expression vectors for human estrogen receptor α , β or androgen receptor (Takeuchi et al., 2005).

Conclusion

Effects on Fertility

Data were insufficient to determine fertility effects of DnOP in humans.

In animals, no effects on number of litters, litter size, sex ratios, pup weight or viability were observed for DnOP in an oral continuous breeding reproductive/developmental study in mice. The NOAEL for fertility effects was 7500 mg/kg bw/d.

In a 13-week subchronic study in rats, no effects on testes weight or morphology were observed at the highest dose level (403 mg/kg bw/d). No testicular atrophy or histological lesions were observed in male rats gavaged with DnOP at 2800 mg/kg bw/d for 4 or 10 days. Decreased testicular zinc was the only effect observed in a one-week rat study in which rats were fed a diet containing 2% DnOP. No effects on testes weights or testicular concentration of testosterone or dihydrotestosterone were observed.

Developmental Effects

Decreased foetal weight and increased visceral malformations were noted in the offspring of rats given high doses of DnOP (up to 9780 mg/kg bw/d) by intraperitoneal injection. However, potentially confounding maternal effects were not reported. In a second short-term oral (gavage) study in mice, statistically significant decreases were observed in viability and pup weight gains. However, concurrent control values for these parameters were not comparable with those of other control groups in the study casting uncertainties over the biological significance of these findings. In contrast, no developmental toxicity was observed in a continuous breeding multigenerational reproductive/developmental study in mice at concentrations up to 7500 mg/kg bw/d.

In view of the magnitude of doses used, the results of two oral studies in mice suggest a low potential for DnOP to induce developmental effects. The NOAEL for developmental effects was 7500 mg/kg bw/d.

4. HAZARD CHARACTERISATION

A full set of toxicity data for DnOP was not available for all health endpoints. For endpoints with missing or incomplete data, information from structurally similar phthalates, where available, was used to extrapolate potential toxicity. Relevant read-across information was obtained from other NICNAS assessment reports for relevant phthalates and the NICNAS Phthalates Hazard Compendium (2007), which contains a comparative analysis of toxicity endpoints across 25 phthalates, including DnOP.

DnOP is a linear C8 phthalate and a member of the High Molecular Weight Phthalate Esters (HMWPEs) Category as defined by the Phthalate Esters Panel HPV Testing Group (2001) and OECD (2004). The HMWPE group includes chemically similar substances produced from alcohols having backbone carbon lengths of $\geq C7$. Due to their similar chemical structure, category members are generally similar with respect to physicochemical, biological and toxicological properties or display an expected trend. Thus, read-across for toxicity endpoints is an appropriate approach to characterise selected endpoints for members of this category.

DnOP is rapidly absorbed from the GIT system following oral administration and metabolised predominantly to mono-n-octylphthalate. In rats after oral administration, peak blood and testes levels of mono-n-octylphthalate are reached at 3 and 6 hours respectively. Elimination occurs via the urine, with levels of MnOP exceeded after 24 hours by MCPP.

In animals, DnOP exhibits low acute oral toxicity. Information on dermal or inhalation toxicity is incomplete or unavailable. Data for other high molecular weight phthalates (NICNAS, 2007) suggests that dermal and inhalation toxicity would be expected to be low.

DnOP caused minimal skin and eye irritation in animals. Data are insufficient to determine the sensitisation potential of DnOP. However, it should be noted that phthalates, in general, have low skin sensitisation potential (NICNAS, 2007).

The liver appears to be the primary target organ from repeated exposure to DnOP. Liver toxicity (weight, histological or clinical chemistry changes) caused by DnOP was observed in several repeat dose studies. DnOP also induced ultrastructural changes in the thyroid and kidney. Overall, liver effects from DnOP did not appear to be associated with peroxisome proliferation. From a 13-week dietary study in rats, a LOAEL of 350 mg/kg bw/d was established based on histological changes in the liver and thyroid. The NOAEL was 37 mg/kg bw/d.

DnOP was negative in bacterial mutation and direct DNA damage assays. Mixtures containing DnOP tested negative in bacterial and mammalian mutation assays. Based on available *in vitro* data, DnOP is considered non-genotoxic.

Limited data on carcinogenic potential indicated that DnOP may act as a promoter of pre-neoplastic hepatic lesions in the rat via a non-peroxisome proliferative mechanism. A large increase in numbers of gamma-glutamyltransferase-positive foci was observed in the livers of rats following 26 weeks dietary administration. Gamma-glutamyltransferase is often significantly increased in human tumours and its role in tumour progression and invasion has been suggested. In another study, liver nodules were reported in rats after dietary administration of DnOP for 15 months. Current literature indicates that rodents compared to

humans are particularly susceptible to peroxisome proliferative effects and resultant carcinogenicity (NICNAS, 2007). However, peroxisome proliferation was not a notable effect in available repeated dose toxicity studies of DnOP. Overall, data are insufficient to determine the carcinogenic potential of DnOP.

DnOP did not appear to induce reproductive toxicity at the highest dose tested of 7500 mg/kg bw/d in a continuous breeding study in mice or 403 mg/kg bw/d in a 13-week subchronic study in rats. *In vitro*, a potential for testicular effects is suggested from findings of mono-n-octyl phthalate induced marked germ-cell detachment and disruption of Sertoli-cell monolayers in co-cultures of Sertoli-germ cells isolated from pubertal rats. However, testicular effects were not seen *in vivo*. The NOAEL for fertility effects was 7500 mg/kg bw/d.

Decreased foetal weight and increased visceral malformations were noted in the offspring of rats given high doses of DnOP post-gestation by intraperitoneal injection. However, the extent to which these might be attributable to maternal toxicity is not known. In a second short-term developmental study in mice, statistically significant decreases were observed in viability and pup weight gains with high doses of DnOP given via oral gavage. However, there are uncertainties over the biological significance of these findings given the magnitude of concurrent control values. In contrast, no developmental toxicity was observed in a continuous breeding multigenerational reproductive/developmental study in mice at DnOP concentrations up to 7500 mg/kg bw/d. In studies of potential estrogenic mimicry, DnOP did not induce estrogenic responses *in vivo* or *in vitro* and displayed no binding affinity for human estrogen receptors.

Studies in mice suggest a low potential for DnOP to induce developmental effects with a NOAEL for developmental effects of 7500 mg/kg bw/d.

The critical endpoints for DnOP are liver and thyroid effects. For these, a LOAEL of 350 mg/kg bw/d was established based on histological changes in these organs. The NOAEL was 37 mg/kg bw/d.

5. HUMAN HEALTH HAZARD SUMMARY TABLE

<i>Phthalate</i>	<i>Acute Toxicity</i>	<i>Irritation & Sensitisation</i>	<i>Repeated Dose Toxicity</i>	<i>Genetic Toxicity</i>	<i>Carcinogenicity</i>	<i>Fertility</i>	<i>Developmental Toxicity</i>
Di-n-octyl phthalate (DnOP)	<p>Oral Rat: LD50 = 53700 mg/kg bw</p> <p>Dermal Guinea pig: LD50 = 75 ml/kg bw</p> <p>Inhalation: no data</p>	<p>Skin Irritation: ME</p> <p>Eye Irritation: ME</p> <p>Respiratory irritation: no data</p> <p>Skin Sensitisation: negative</p>	<p>Oral Rat: NOAEL = 37 mg/kg bw/d</p> <p>LOAEL = 350 mg/kg bw/d: ultrastructural changes in liver and thyroid.</p> <p>High doses: ↑ liver weight, vacuolization and necrosis; ↓ centrilobular glycogen. PP not generally noted.</p>	<p><i>In vitro</i> Bacterial mutation assays and direct DNA damage assays: negative</p> <p><i>In vivo</i> No data.</p>	<p><i>In vivo</i> Rat: Liver nodules following 15 month dietary administration.</p> <p>Rat: 26 week dietary ↑ gamma-glutamyltransferase-positive foci in liver</p> <p><i>In vitro</i> No data.</p>	<p>Continuous breeding study Mouse: NOAEL = 7500 mg/kg bw/d (highest dose tested)</p> <p>LOAEL: NE</p>	<p>Continuous breeding study Mouse: NOAEL = 7500 mg/kg bw/d (highest dose tested)</p> <p>LOAEL: NE</p>

ME – minimal effects; NE – not established; PP – peroxisome proliferation

6. REFERENCES

- Albro PW and Moore B (1974) Identification of the metabolites of simple phthalate diesters in rat urine. *J. Chromatography* 94: 209-218
- Anon. (1946) *American J. Ophthalmology Series 3: V.1-1918* (29,1363, 1946).
- ATSDR (1997) Toxicological Profile for Di-*n*-Octylphthalate. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, September 1997
- Barber ED, Cifone M, Rundell J, Przygoda R, Astill BD, Moran E, Mulholland A, Robinson E & Schneider B (2000) Results of the L5178Y mouse lymphoma assay and the Balb/3T3 cell in vitro transformation assay for eight phthalate esters. *J. Appl. Toxicol.* 20:69-80.
- Bisesi MS (1994) Esters. *In: Clayton GD and Clayton FE (Eds.) Patty's Industrial Hygiene and Toxicology* 4th ed. John Wiley and Sons, Inc. New York. 1994
- Calafat AM, Silva MJ, Reidy JA, Gray LE, Samandar E, Preau JL, Herbert AR, Needham LL (2006) Mono-(3-carboxypropyl) phthalate, a metabolite of di-*n*-octyl phthalate. *J. Toxicol. Environ. Health.* 69:215-227
- Carter JH, Richmond RE, Carter HW, et al. (1992) Quantitative image cytometry of hepatocytes expressing gamma-glutamyl transpeptidase and glutathione S-transferase in diethylnitrosamine-initiated rats treated with phenobarbital and/or phthalate esters. *J Histochem Cytochem* 40: 1105-1115.
- Carter JH, Carter HW, DeAngelo AB & Daniel FB (1989) Sub-lethal Autolysis in Livers of Rats Exposed to Phthalates. *J. Cell Biol.* 109(4): 182a, Abstract No. 1003.
- CERHR (2003) NTP-CERHR Monograph on the Potential Human Reproductive and Developmental Effects of Di-*n*-Octyl Phthalate (DnOP) CERHR NIH Publication No. 03-4488 (May 2003). National Toxicology Program, US Department of Health and Human Services.
- CMA (1999) Comments of the Chemical Manufacturers Association phthalate esters panel in response to request for public input on seven phthalate esters. FR Doc. 99-9484. Washington, DC: Chemical Manufacturers Association.
- DeAngelo AB, Garrett CT, & Daniel FB (1989) Di-*n*-octyl phthalate but not di(2-ethylhexyl)phthalate promotes diethylnitrosamine initiated hepatocellular carcinoma in the male F344 rat. *Proceedings of the American Association of Cancer Research* 30:204.
- DeAngelo AB, Garrett CT, & Manolukas, LA, et al. (1986) Di-*n*-octyl phthalate (DOP), a relatively ineffective peroxisome inducing straight chain isomer of the environmental contaminant di(2-ethylhexyl)phthalate (DEHP), enhances the development of putative preneoplastic lesions in rat liver. *Toxicology* 41:279-288.
- DeAngelo AB, Cicmanec J, McMillan LP, & Wernsing PA (1988) Comparative Toxicity of Di(2-ethylhexyl) Phthalate (DEHP) and Di-*n*-octyl Phthalate (DOP). *The Toxicologist* 8(1): 38, Abstract No. 150.
- Dogra RK, Chandra K, Chandra S et al (1989) Di-octyl phthalate induced altered host resistance: Viral and protozoal models in mice. *Industrial Health* 27: 83-87
- Dogra RK, Khanna S, Shukla L et al (1987) Modification of the immune response in rats by di-octyl phthalate. *Industrial Health* 25: 97-101

- Eastman Kodak Company (1978) Toxicity and health hazard summary. Rochester, NY: EPA/OTS Document No. 878214345
- ECB (European Chemicals Bureau) (2000) IUCLID Dataset Existing Chemical Substance ID: 71662-46-9. European Commission.
- Foster PMD, Thomas LV, Cook MW, Gangolli SD (1980) Study of the testicular effects and changes in zinc excretion produced by some n-alkyl phthalates in the rat. *Toxicol Appl Pharmacol* 54:392-398.
- Fredricsson B, Muller L, Pousette A, & Westerholm R (1993) Human sperm motility is affected by plasticisers and diesel particle extracts. *Pharm Tox* 72: 128-133.
- Gilioli R, Bulgheroni C, Terrano T, Filippini G, Massetto G & Boeri R (1978) Studio neurologico trasversale e longitudinale di una popolazione operaia addetta alla produzione di ftalati. *Med Lav* 69:620-631.
- Goodyear Tire & Rubber Company (1981) DNA damage by dioctyl phthalate BASF, Tank 28 in the E. coli Pol A - Assay. Goodyear Tire and Rubber Company Laboratory Report No. 81-42-2. Akron, OH.
- Government of Canada (1993) Canadian Environmental Protection Act. Priority Substances List Assessment Report. Di-*n*-Octyl Phthalate.
- Gray TJ and Butterworth KR (1980) Testicular atrophy produced by phthalate esters. *Arch. Toxicol. Suppl.* 4: 452-455.
- Gray TJ and Beamand JA (1984) Effects of some phthalate esters and other testicular toxins on primary cultures of testicular cells. *Food Chem Toxicol*, 22:123-131.
- Gulati DK, Chambers R, Shaver S, Sabehrwal PS, & Lamb JC (1985) Di-n-octyl phthalate reproductive and fertility assessment in CD-1 mice when administered in feed. Research Triangle Park: US National Toxicology Program.
- Hardin BD, Schuler RL, Burg JR, Booth GM, Hazelden KP, MacKenzie KM, Piccirillo VJ, & Smith KN (1987) Evaluation of 60 chemicals in a preliminary developmental toxicity test. *Teratogen Carcinogen Mutagen* 7:29-48.
- Harris CA, Henttu P, Parker MG, & Sumpter JP (1997) The estrogenic activity of phthalate esters in vitro. *Environ. Health Perspect* 105: 802-811
- Heindel JJ, Gulati DK, Mounce RC, Russell SR & Lamb JCI (1989) Reproductive toxicity of three phthalic acid esters in a continuous breeding protocol. *Fundam Appl Toxicol* 12:508-518.
- Hellwig J & Jackh R (1997) Differential prenatal toxicity of one straight-chain and five branched chain primary alcohols in rats. *Food Chem Toxicol* 35:489-500.
- Hinton RH, Mitchell FE, Mann A, Chescoe D, Price SC, Nunn A, Grasso P & Bridges JW (1986) Effects of Phthalic Acid Esters on the Liver and Thyroid. *Environ Health Perspect* 70:195-210
- Khanna S, Dogra RKS, Bhatnagar MC, Shukla LJ, Srivastava SN, & Shanker R (1990) Nephrotoxicity of di-octyl phthalate treated rats - histological evidence. *J Environ Biol* 11:27-34.
- Lake BG, Gray TJ, & Gangolli SD (1986) Hepatic effects of phthalate esters and related compounds - in vivo and in vitro correlations. *Environ Health Perspect* 67:283-290.

- Lake BG, Rijcken WR, Gray TJ, et al. (1984) Comparative studies of the hepatic effects of di- and mono-n-octyl phthalates, di-(2-ethylhexyl) phthalate and clofibrate in the rat. *Acta Pharmacol Toxicol* 54:167-176.
- Mann AH, Price SC, Mitchell FE, Grasso P, Hinton R, & Bridge J (1985) Comparison of the short-term effects of di(2-ethylhexyl)phthalate, di(n-hexyl)phthalate, and di(n-octyl)phthalate in rats. *Toxicol Appl Pharmacol* 77:116-132.
- Marhold J (1986) *Prehled Prumyslove Toxikologie; Organicke Latky*. Prague, Czechoslovakia, Avicenum, 1986 (390, 1986).
- Milkov LB, Aldjreva MV, Popova TB et al. (1973) Health Status of Workers Exposed to Effect of Phthalate Plasticizers in the Production of Artificial Leather and Films (on the Basis of PVC Resins). Translation of *Gig Tr Prof Zabol* 13:14-17. NTIS PB-221 973T, p 5.
- Morrissey RE, Lamb JC, & Morris RW, et al. (1989) Results and evaluations of 48 continuous breeding reproduction studies conducted in mice. *Fundam Appl Toxicol* 13:747-777.
- NICNAS (2007) *Phthalates Hazard Compendium*. A summary of physicochemical and human health hazard data for 25 phthalate chemicals. Sydney, National Industrial Chemicals Notification and Assessment Scheme.
- NJDHSS (New Jersey Department of Health and Senior Services) (2002) *Hazardous Substance Fact Sheet*. Di-n-octyl Phthalate.
- NTP (1985) *Di-n-octylphthalate: reproduction and fertility assessment in CD-1 mice when administered in feed*. Research Triangle Park NC: National Institute of Environmental Health Sciences, National Toxicology Program. NTP-85-140. NTIS Order No. PB85-218147
- OECD (2004) *SIDS Initial Assessment Report for SIAM 19: Category – High Molecular Weight Phthalate Esters*. Organisation for Economic Cooperation and Development, Berlin, Germany, 19-22 October 2004.
- Oishi S (1990) Effects of phthalic acid esters on testicular mitochondrial functions in the rat. *Arch. Toxicol.* 64:143-147
- Oishi S, Hiraga K (1980) Testicular atrophy induced by phthalic acid esters: Effect on testosterone and zinc concentrations. *Toxicol Appl Pharmacol* 53:35-41.
- Oishi S & Hiraga K (1982) Effects of monoesters of O-phthalic acid on serum lipid composition of rats. *Toxicol Lett* 14:79-84
- Ontario Ministry of the Environment (2005) *Ontario Air Standards For Di-n-Octyl Phthalate (DNOP)*. Ontario, Canada. June 2005
- Phthalate Esters Panel HPV Testing Group (2001) *High production volume (HPV) chemical challenge programme test plan for the phthalate esters category*. December 10, 2001.
- Pieckacz H (1971) The influence on rat organism of dibutyl phthalate and dioctyl phthalate administered perorally for a prolonged period. III. The influence on the fertility and foetal development. *Rocz. Panstw. Zakl. Hig.* 22(5): 519-526 (Polish, with English abstract).
- Poon R, Lecavalier P, Mueller R, et al. (1995) *Subchronic oral toxicity of di-n-octylphthalate and di(2-ethylhexyl)phthalate in the rat*. [ATSDR peer-reviewed draft].

- Poon R, Lecavalier P, Mueller R, Valli VE, Proctor BG, & Chu I (1997) Subchronic oral toxicity of di-n-octyl phthalate and di(2-ethylhexyl)phthalate in the rat. *Food and Chem Toxicol* 35:225-239.
- Reddy BS, Rozati R, Reddy BVR and Raman NVVSS (2006a) Association of phthalate esters with endometriosis in Indian women. *BJOG: An International Journal of Obstetrics & Gynaecology* 113(5):515-520.
- Reddy BS, Rozati R, Reddy S, Kodampur S, Reddy, P and Reddy, R (2006b) High plasma concentrations of polychlorinated biphenyls and phthalate esters in women with endometriosis: a prospective case control study. *Fertility & Sterility* 85: 775-779.
- Rowland IR, Cottrel RC and Phillips JC (1977)* Hydrolysis of phthalate esters by the gastrointestinal contents of the rat. *Food Cosmet. Toxicol.* 15:17-21
- RTECS (2004) Phthalic acid, dioctyl ester. Registry of toxic effects of chemical substances. US National Institute of Occupational Safety and Health.
- Sato T, Nagase H, Sato K, et al. (1994) Enhancement of the mutagenicity of amino acid pyrolysates by phthalate esters. *Environ Mol Mut* 24:325-331.
- Seed JL (1982) Mutagenic activity of phthalate esters in bacterial liquid suspension assays. *Environ. Health Perspect.* 45: 111-114
- Shibamoto T and Wei CI (1986) Mutagenicity of materials extracted from synthetic rubber. *Agric. Biol. Chem.* 50: 513-514
- Silva MJ, Kato K, Gray EL, Wolf C, Heedham LL and Calafat AM (2005) Urinary metabolites of di-n-octyl phthalate in rats. *Toxicology* 210: 123-133
- Singh AR, Lawrence WH, & Autian J (1972) Teratogenicity of phthalate esters in rats. *J Pharm Sci*, 61:51-55.
- Swan S, Main K, Liu F, Stewart S, Kruse R, Calafat A, Mao C, Redmon J, Ternand C, Sullivan S, Teague J, & The-Study-for-Future-Families-Research-Team (2005) Decrease in anogenital distance among male infants with prenatal phthalate exposure. *Environ Health Perspect* 113: 1056-1061.
- Takeuchi S, Iida M, Kobayashi S, Jin K, Matsuda T, & Kojima H (2005) Differential effects of phthalate esters on transcriptional activities via human estrogen receptors alpha and beta and androgen receptor. *Toxicology* 210: 223-233.
- Toda C, Okamoto Y, Ueda K, Hashizume K, Itoh K, & Kojima N (2004) Unequivocal estrogen receptor-binding affinity of phthalate esters featured with ring hydroxylation and proper alkyl chain size. *Arch Biochem Biophys* 431: 16-21.
- Zacharewski TR, Meek MD, Clemons JH, Wu ZF, Fielden MR, & Matthews JB (1998) Examination of the in vitro and in vivo estrogenic activities of eight commercial phthalate esters. *Toxicol Sci* 46:282-293.
- Zeiger E, Haworth S, Mortelmans K, et al. (1985) Mutagenicity testing of di(Zethylhexyl)phthalate and related chemicals in salmonella. *Environ Mutagen* 7:213-232.
- Zeiger E, Haworth S, Speck W, et al. (1982) Phthalate ester testing in the National Toxicology Program's environmental mutagenesis test development program. *Environ Health Perspect* 45:99-101